

Fluoxetine

Core Safety Profile

4.3 Contraindications

Hypersensitivity to fluoxetine or to any of its excipients.

Monoamine Oxidase Inhibitors: Cases of serious and sometimes fatal reactions have been reported in patients receiving an SSRI in combination with a monoamine oxidase inhibitor (MAOI), and in patients who have recently discontinued an SSRI and have been started on a MAOI. Treatment of fluoxetine should only be started 2 weeks after discontinuation of an irreversible MAOI and the following day after discontinuation of a reversible MAOI-A.

Some cases presented with features resembling serotonin syndrome (which may resemble and be diagnosed as neuroleptic malignant syndrome). Cyproheptadine or dantrolene may benefit patients experiencing such reactions. Symptoms of a drug interaction with a MAOI include: hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, mental status changes that include confusion, irritability and extreme agitation progressing to delirium and coma.

Therefore, fluoxetine is contra-indicated in combination with a non-selective MAOI. Similarly, at least 5 weeks should elapse after discontinuing fluoxetine treatment before starting a MAOI. If fluoxetine has been prescribed chronically and/or at a high dose, a longer interval should be considered.

The combination of fluoxetine with a reversible MAOI (e.g. moclobemide) is not recommended. Treatment with fluoxetine can be initiated the following day after discontinuation of a reversible MAOI.

4.4. Special warnings and precautions for use

Use in children and adolescents under 18 years of age

Suicide-related behaviours (suicide attempt and suicidal thoughts), and hostility (predominantly aggression, oppositional behaviour and anger) were more frequently observed in clinical trials among children and adolescents treated with antidepressants compared to those treated with placebo. Prozac should only be used in children and adolescents aged 8 to 18 years for the treatment of moderate to severe major depressive episodes and it should not be used in other indications. If, based on clinical need, a decision to treat is nevertheless taken, the patient should be carefully monitored for the appearance of suicidal symptoms. In addition, only limited evidence is available concerning long-term effect on safety in children and adolescents, including effects on growth, sexual maturation and cognitive, emotional and behavioural developments (see section 5.3).

In a 19-week clinical trial decreased height and weight gain was observed in children and adolescents treated with fluoxetine (see section 4.8). It has not been established whether there is an effect on achieving normal adult height. The possibility of a delay in puberty cannot be ruled out (see sections 5.3 and 4.8). Growth and pubertal development (height, weight and TANNER staging) should therefore be monitored during and after treatment with fluoxetine. If either is slowed, referral to a paediatrician should be considered.

In paediatric trials, mania and hypomania were commonly reported (see section 4.8). Therefore, regular monitoring for the occurrence of mania/hypomania is recommended. Fluoxetine should be discontinued in any patient entering a manic phase.

It is important that the prescribers discusses carefully the risks and benefits of treatment with the child/young person and/or their parents.

Rash and allergic reactions: Rash, anaphylactoid events and progressive systemic events, sometimes serious (involving skin, kidney, liver or lung) have been reported. Upon the appearance of rash or of other allergic phenomena for which an alternative aetiology cannot be identified, fluoxetine should be discontinued.

Seizures: Seizures are a potential risk with antidepressant drugs. Therefore, as with other antidepressants, fluoxetine should be introduced cautiously in patients who have a history of seizures. Treatment should be discontinued in any patient who develops seizures or where there is an increase in seizure frequency. Fluoxetine should be avoided in patients with unstable seizure disorders/epilepsy and patients with controlled epilepsy should be carefully monitored.

Mania: Antidepressants should be used with caution in patients with a history of mania/hypomania. As with all antidepressants, fluoxetine should be discontinued in any patient entering a manic phase.

Hepatic/Renal Function: Fluoxetine is extensively metabolised by the liver and excreted by the kidneys. A lower dose, e.g., alternate day dosing, is recommended in patients with significant hepatic dysfunction. When given fluoxetine 20 mg/day for 2 months, patients with severe renal failure (GFR <10 ml/min) requiring dialysis showed no difference in plasma levels of fluoxetine or norfluoxetine compared to controls with normal renal function.

Cardiac Disease: No conduction abnormalities that resulted in heart block were observed in the ECG of 312 patients who received fluoxetine in double blind clinical trials. However, clinical experience in acute cardiac disease is limited, therefore caution is advisable.

Weight Loss: Weight loss may occur in patients taking fluoxetine but it is usually proportional to baseline body weight.

Diabetes: In patients with diabetes, treatment with an SSRI may alter glycaemic control. Hypoglycaemia has occurred during therapy with fluoxetine and hyperglycaemia has developed following discontinuation. Insulin and/or oral hypoglycaemic dosage may need to be adjusted.

Suicide/suicidal thoughts or clinical worsening: Depression is associated with an increased risk of suicidal thoughts, self harm and suicide (suicide-related events). This risk persists until significant remission occurs. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored until such improvement occurs. It is general clinical experience that the risk of suicide may increase in the early stages of recovery.

Other psychiatric conditions for which Prozac is prescribed can also be associated with an increased risk of suicide-related events. In addition, these conditions may be co-morbid with major depressive disorder. The same precautions observed when treating patients with major depressive disorder should therefore be observed when treating patients with other psychiatric disorders.

Patients with a history of suicide-related events, those exhibiting a significant degree of suicidal ideation prior to commencement of treatment are known to be at greater risk of suicidal thoughts or suicide attempts, and should receive careful monitoring during treatment. A meta-analysis of placebo-controlled clinical trials of antidepressants drugs in adult patients with psychiatric disorders showed an increased risk of suicidal behaviour with antidepressants compared to placebo in patients less than 25 years old.

Close supervision of patients and in particular those at high risk should accompany drug therapy especially in early treatment and following dose changes. Patients (and caregivers of patients) should be alerted about the need to monitor for any clinical worsening, suicidal behaviour or thoughts and unusual changes in behaviour and to seek medical advice immediately if these symptoms present.

Akathisia/psychomotor restlessness: The use of fluoxetine has been associated with the development of akathisia, characterised by a subjectively unpleasant or distressing restlessness and need to move often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.

Withdrawal symptoms seen on discontinuation of SSRI treatment: Withdrawal symptoms when treatment is discontinued are common, particularly if discontinuation is abrupt (see section 4.8). In clinical trials adverse events seen on treatment discontinuation occurred in approximately 60% of patients in both the fluoxetine and placebo groups. Of these adverse events, 17% in the fluoxetine group and 12% in the placebo group were severe in nature.

The risk of withdrawal symptoms may be dependent on several factors including the duration and dose of therapy and the rate of dose reduction. Dizziness, sensory disturbances (including paraesthesia), sleep disturbances (including insomnia and intense dreams), asthenia, agitation or anxiety, nausea and/or vomiting, tremor and headache are the most commonly reported reactions. Generally these symptoms are mild to moderate however, in some patients they may be severe in intensity. They usually occur within the first few days of discontinuing treatment. Generally these symptoms are self-limiting and usually resolve within 2 weeks, though in some individuals they may be prolonged (2-3 months or more). It is therefore advised that Prozac should be gradually tapered when discontinuing treatment over a period of at least one to two weeks,

according to the patient's needs (see *Withdrawal symptoms seen on discontinuation of Prozac*, section 4.2).

Haemorrhage: There have been reports of cutaneous bleeding abnormalities such as ecchymosis and purpura with SSRI's. Ecchymosis has been reported as an infrequent event during treatment with fluoxetine. Other hemorrhagic manifestations (e.g., gynaecological haemorrhages, gastrointestinal bleedings and other cutaneous or mucous bleedings) have been reported rarely. Caution is advised in patients taking SSRI's, particularly in concomitant use with oral anticoagulants, drugs known to affect platelet function (e.g. atypical antipsychotics such as clozapine, phenothiazines, most TCA's, aspirin, NSAID's) or other drugs that may increase risk of bleeding as well as in patients with a history of bleeding disorders.

Electroconvulsive Therapy (ECT): There have been rare reports of prolonged seizures in patients on fluoxetine receiving ECT treatment, therefore caution is advisable.

St John's Wort: An increase in serotonergic effects, such as serotonin syndrome, may occur when selective serotonin reuptake inhibitors and herbal preparations containing St John's Wort (*Hypericum perforatum*) are used together.

On rare occasions development of a serotonin syndrome or neuroleptic malignant syndrome-like events have been reported in association with treatment of fluoxetine, particularly when given in combination with other serotonergic (among others L-tryptophan) and/or neuroleptic drugs. As these syndromes may result in potentially life-threatening conditions, treatment with fluoxetine should be discontinued if such events (characterised by clusters of symptoms such as hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, mental status changes including confusion, irritability, extreme agitation progressing to delirium and coma) occur and supportive symptomatic treatment should be initiated.

4.5. Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

Half-life: The long elimination half-lives of both fluoxetine and norfluoxetine should be borne in mind (see section 5.2) when considering pharmacodynamic or pharmacokinetic drug interactions (e.g. when switching from fluoxetine to other antidepressants).

Monoamine oxidase inhibitors: (see section 4.3).

Not recommended combinations: MAOI-A (see section 4.3)

Combinations requiring precautions for use: MAOI-B (selegiline): risk of serotonin syndrome. Clinical monitoring is recommended.

Phenytoin: Changes in blood levels have been observed when combined with fluoxetine. In some cases manifestations of toxicity have occurred. Consideration should be given to using conservative titration schedules of the concomitant drug and to monitoring clinical status.

Serotonergic drugs: Co-administration with serotonergic drugs (e.g. tramadol, triptans) may increase the risk of serotonin syndrome. Use with triptans carries the additional risk of coronary vasoconstriction and hypertension.

Lithium and tryptophan: There have been reports of serotonin syndrome when SSRIs have been given with lithium or tryptophan and, therefore, the concomitant use of fluoxetine with these drugs should be undertaken with caution. When fluoxetine is used in combination with lithium, closer and more frequent clinical monitoring is required.

CYP2D6 isoenzyme: Because fluoxetine's metabolism (like tricyclic antidepressants and other selective serotonin antidepressants) involves the hepatic cytochrome CYP2D6 isoenzyme system, concomitant therapy with drugs also metabolised by this enzyme system may lead to drug interactions. Concomitant therapy with drugs predominantly metabolised by this isoenzyme, and which have a narrow therapeutic index (such as flecainide, encainide, carbamazepine and tricyclic antidepressants), should be initiated at or adjusted to the low end of their dose range. This will also apply if fluoxetine has been taken in the previous 5 weeks.

Oral anticoagulants: Altered anti-coagulant effects (laboratory values and/or clinical signs and symptoms), with no consistent pattern, but including increased bleeding, have been reported uncommonly when fluoxetine is co-administered with oral anticoagulants. Patients receiving warfarin therapy should receive careful coagulation monitoring when fluoxetine is initiated or stopped. (see section 4.4, *Haemorrhage*).

Electroconvulsive Therapy (ECT): There have been rare reports of prolonged seizures in patients on fluoxetine receiving ECT treatment, therefore caution is advisable.

Alcohol: In formal testing, fluoxetine did not raise blood alcohol levels or enhance the effects of alcohol. However, the combination of SSRI treatment and alcohol is not advisable.

St. John's Wort: In common with other SSRIs, pharmacodynamic interactions between fluoxetine and the herbal remedy St. John's Wort (*Hypericum perforatum*) may occur, which may result in an increase of undesirable effects.

4.6. Pregnancy and lactation

Pregnancy:

Some epidemiological studies suggest an increased risk of cardiovascular defects associated with the use of fluoxetine during the first trimester. The mechanism is unknown. Overall the data suggest that the risk of having an infant with a cardiovascular defect following maternal fluoxetine exposure is in the region of

2/100 compared with an expected rate for such defects of approximately 1/100 in the general population.

Epidemiological data have suggested that the use of SSRIs in pregnancy, particular in late pregnancy, may increase the risk of persistent pulmonary hypertension in the newborn (PPHN). The observed risk was approximately 5 cases per 1000 pregnancies. In the general population 1 to 2 cases of PPHN per 1000 pregnancies occur.

Furthermore, although fluoxetine can be used during pregnancy, caution should be exercised, especially during late pregnancy or just prior to the onset of labour since some other effects have been reported in neonates: irritability, tremor, hypotonia, persistent crying, difficulty in sucking or in sleeping. These symptoms may indicate either serotonergic effects or a withdrawal syndrome. The time to occur and the duration of these symptoms may be related to the long half-life of fluoxetine (4-6 days) and its active metabolite, norfluoxetine (4-16 days).

Lactation: Fluoxetine and its metabolite norfluoxetine, are known to be excreted in human breast milk. Adverse events have been reported in breastfeeding infants. If treatment with fluoxetine is considered necessary, discontinuation of breastfeeding should be considered; however, if breastfeeding is continued, the lowest effective dose of fluoxetine should be prescribed.

4.7. Effects on ability to drive and use machines

Although fluoxetine has been shown not to affect psychomotor performance in healthy volunteers, any psychoactive drug may impair judgement or skills. Patients should be advised to avoid driving a car or operating hazardous machinery until they are reasonably certain that their performance is not affected.

4.8. Undesirable effects

Undesirable effects may decrease in intensity and frequency with continued treatment and do not generally lead to cessation of therapy.

In common with other SSRIs the following undesirable effects have been seen:

Body as a whole: Hypersensitivity (e.g. pruritis, rash, urticaria, anaphylactoid reaction, vasculitis, serum sickness-like reaction, angioedema) (see sections 4.3 and 4.4), chills, serotonin syndrome, photosensitivity, and very rarely Erythema Multiforme that could progress to Stevens-Johnson syndrome or Toxic Epidermal Necrolysis (Lyell syndrome).

Digestive system: Gastrointestinal disorders (e.g. diarrhoea, nausea, vomiting, dyspepsia, dysphagia, taste perversion), dry mouth. Abnormal liver function tests have been reported rarely. Very rare cases of idiosyncratic hepatitis.

Nervous system: Headache, sleep abnormalities (e.g. abnormal dreams, insomnia), dizziness, anorexia, fatigue (e.g. somnolence, drowsiness), euphoria, transient abnormal movement (e.g., twitching, ataxia, tremor, myoclonus), seizures and rarely psychomotor restlessness/akathisia (see section 4.4). Hallucinations, manic reaction, confusion, agitation, anxiety and associated symptoms (e.g. nervousness), impaired concentration and thought process (e.g. depersonalisation), panic attacks, very rarely serotonin syndrome, suicidal thoughts and behaviour (these symptoms may be due to the underlying disease). Cases of suicidal ideation and suicidal behaviour have been reported during fluoxetine therapy or early after treatment discontinuation (see Section 4.4)

Urogenital system: Urinary retention, urinary frequency

Reproductive disorders: Sexual dysfunction (delayed or absent ejaculation, anorgasmia), priapism, galactorrhoea.

Miscellaneous: Alopecia, yawn, abnormal vision (e.g., blurred vision, mydriasis), sweating, vasodilatation, arthralgia, myalgia, postural hypotension, ecchymosis. Other haemorrhagic manifestations (e.g., gynaecological haemorrhages, gastrointestinal bleedings and other cutaneous or mucous bleedings) have been reported rarely (see section 4.4, *Haemorrhage*).

Hyponatraemia: Hyponatraemia (including serum sodium below 110 mmol/l) has been rarely reported and appeared to be reversible when fluoxetine was discontinued. Some cases were possibly due to the syndrome of inappropriate antidiuretic hormone secretion. The majority of reports were associated with older patients, and patients taking diuretics or otherwise volume depleted.

Respiratory system: Pharyngitis, dyspnoea. Pulmonary events (including inflammatory processes of varying histopathology and/or fibrosis) have been reported rarely. Dyspnoea may be the only preceding symptom.

Bone fractures: Epidemiological studies, mainly conducted in patients 50 years of age and older, show an increased risk of bone fractures in patients receiving SSRIs and TCAs. The mechanism leading to the risk is unknown.

Withdrawal symptoms seen on discontinuation of fluoxetine treatments: Discontinuation of fluoxetine commonly leads to withdrawal symptoms. Dizziness, sensory disturbances (including paraesthesia), sleep disturbances (including insomnia and intense dreams), asthenia, agitation or anxiety, nausea and/or vomiting, tremor and headache are the most commonly reported reactions. Generally these events are mild to moderate and are self-limiting, however, in some patients they may be severe and/or prolonged (see section 4.4). It is therefore advised that when Prozac treatment is no longer required, gradual discontinuation by dose tapering should be carried out (see sections 4.2 and 4.4).

Children and adolescents (see section 4.4):

In paediatric clinical trials suicide-related behaviours (suicide attempt and suicidal thoughts), and hostility were more frequently observed among children and adolescents treated with antidepressants compared to those treated with placebo.

The safety of fluoxetine has not been systematically assessed for chronic treatment longer than 19 weeks.

In paediatric clinical trials, manic reactions, including mania and hypomania, were reported (2.6% of fluoxetine-treated patients vs. 0% in placebo-controls), leading to discontinuation in the majority of cases. These patients had no prior episodes of hypomania/mania.

After 19 weeks of treatment, paediatric subjects treated with fluoxetine in a clinical trial gained an average of 1.1 cm less in height ($p=0.004$) and 1.1 kg less in weight ($p=0.008$) than subjects treated with placebo. Isolated cases of growth retardation have also been reported from clinical use.

Isolated cases of adverse events potentially indicating delayed sexual maturation or sexual dysfunction have been reported from paediatric clinical use. (see also section 5.3)

In paediatric clinical trials, fluoxetine treatment was associated with a decrease in alkaline phosphatase levels.

4.9. Overdose

Cases of overdose of fluoxetine alone usually have a mild course. Symptoms of overdose have included nausea, vomiting, seizures, cardiovascular dysfunction ranging from asymptomatic arrhythmias to cardiac arrest, pulmonary dysfunction, and signs of altered CNS status ranging from excitation to coma. Fatality attributed to overdose of fluoxetine alone has been extremely rare. Cardiac and vital signs monitoring are recommended, along with general symptomatic and supportive measures. No specific antidote is known.

Forced diuresis, dialysis, haemoperfusion, and exchange transfusion are unlikely to be of benefit. Activated charcoal, which may be used with sorbitol, may be as or more effective than emesis or lavage. In managing overdosage, consider the possibility of multiple drug involvement. An extended time for close medical observation may be needed in patients who have taken excessive quantities of a tricyclic antidepressant if they are also taking, or have recently taken, fluoxetine.